

NOTE ON COFIBRATIONS

ARNE STRØM

In the first section of this note it is proved that cofibrations are homeomorphisms, and a characterization of closed cofibrations is given. The second section contains the proof of a homotopy lifting-extension theorem generalizing a result on relative CW-complexes.

All functions considered will be continuous.

1.

DEFINITION. A (Hurewicz) *fibration* is a map $p: E \rightarrow B$ with the property that for any map $f: X \rightarrow E$ and any homotopy $F: X \times I \rightarrow B$ such that $F(x, 0) = pf(x)$ for all $x \in X$, there exists a homotopy $\bar{F}: X \times I \rightarrow E$ such that $p\bar{F} = F$ and $\bar{F}(x, 0) = f(x)$ for all $x \in X$.

A *cofibration* is a map $j: A \rightarrow X$ such that for any map $f: X \rightarrow Y$ and any homotopy $\bar{F}: A \times I \rightarrow Y$ such that $\bar{F}(a, 0) = fj(a)$ for all $a \in A$, there exists a homotopy $F: X \times I \rightarrow Y$ such that $F(j \times 1_I) = \bar{F}$ and $F(x, 0) = f(x)$ for all $x \in X$.

If A is a subspace of a space X such that the inclusion map $A \subset X$ is a cofibration, the pair (X, A) is called a *cofibered pair* or is said to possess the *absolute homotopy extension property* (AHEP). A necessary condition for (X, A) to be a cofibered pair is the existence of a retraction

$$r: X \times I \rightarrow (X \times 0) \cup (A \times I).$$

If A is closed, this condition is also sufficient.

The following theorem shows that the only cofibrations are cofibered pairs.

THEOREM 1. *If $j: A \rightarrow X$ is a cofibration, then j is a homeomorphism $A \approx j(A)$.*

PROOF. Let $j: A \rightarrow X$ be a cofibration and consider the mapping cylinder Z of j , that is, the quotient space of the topological sum $(X \times 0) \vee (A \times I)$ obtained by identifying $(a, 0) \in A \times I$ with $(j(a), 0) \in X \times 0$

for each $a \in A$. Denote by q the quotient map $(X \times 0) \vee (A \times I) \rightarrow Z$. There is a continuous map $i: Z \rightarrow X \times I$ defined by

$$\begin{aligned} iq(x, 0) &= (x, 0), & x \in X, \\ iq(a, t) &= (j(a), t), & a \in A, t \in I. \end{aligned}$$

Define maps $f: X \rightarrow Z$ and $\bar{F}: A \times I \rightarrow Z$ by

$$f(x) = q(x, 0), \quad \bar{F}(a, t) = q(a, t).$$

Because j is a cofibration there exists a map $F: X \times I \rightarrow Z$ such that $F(j(a), t) = q(a, t)$ and $F(x, 0) = q(x, 0)$ for all $a \in A$, $t \in I$, and $x \in X$. Then $F \circ i = 1_Z$, and i is, therefore, a homeomorphism of Z onto $i(Z) = (X \times 0) \cup (j(A) \times I)$. Also, $q|_{A \times 1}$ is a homeomorphism of $A \times 1$ onto $q(A \times 1)$, and consequently $i q|_{A \times 1}$ is a homeomorphism of $A \times 1$ onto $i q(A \times 1) = j(A) \times 1$.

Next we shall prove a theorem which generalizes 3.1 of [1].

THEOREM 2. *Let A be a closed subspace of a topological space X . Then (X, A) is a cofibered pair if and only if there exist*

- (i) *a neighborhood U of A which is deformable in X to A rel A (that is, there exists a homotopy $H: U \times I \rightarrow X$ such that $H(x, 0) = x$, $H(a, t) = a$, and $H(x, 1) \in A$ for all $x \in U$, $a \in A$, $t \in I$), and*
- (ii) *a continuous function $\varphi: X \rightarrow I$ such that $A = \varphi^{-1}(0)$ and $\varphi(x) = 1$ for all $x \in X - U$.*

PROOF. Suppose that (X, A) is a cofibered pair. Then there exists a retraction

$$r: X \times I \rightarrow (X \times 0) \cup (A \times I),$$

and U , H and φ may be chosen as follows:

$$\begin{aligned} U &= \{x \in X \mid pr_1 r(x, 1) \in A\}, \\ H &= pr_1 r|_{U \times I}, \\ \varphi(x) &= \sup_{t \in I} |t - pr_2 r(x, t)|, \end{aligned}$$

pr_1 and pr_2 denoting projections on X and I , respectively.

Conversely, suppose that U , H and φ are given and satisfy the conditions of the theorem. Since A is closed it suffices to prove the existence of a retraction

$$r: X \times I \rightarrow (X \times 0) \cup (A \times I).$$

The required retraction may be constructed as follows.

- (i) If $\varphi(x) = 1$, let $r(x, t) = (x, 0)$.

- (ii) If $\frac{1}{2} \leq \varphi(x) < 1$, let $r(x, t) = (H(x, 2(1 - \varphi(x))t), 0)$.
 (iii) If $0 < \varphi(x) \leq \frac{1}{2}$ and $0 \leq t \leq 2\varphi(x)$, let

$$r(x, t) = (H(x, t/(2\varphi(x))), 0) .$$

 (iv) If $0 < \varphi(x) \leq \frac{1}{2}$ and $2\varphi(x) \leq t \leq 1$, let

$$r(x, t) = (H(x, 1), t - 2\varphi(x)) .$$

 (v) If $\varphi(x) = 0$, let $r(x, t) = (x, t)$.

(This construction is that of [2].) The proof of continuity is straightforward and will be omitted.

2.

It was remarked in Section 1 that if (X, A) is a cofibered pair, then $(X \times 0) \cup (A \times I)$ is a retract of $X \times I$. In fact, we have the following stronger result.

LEMMA. *If (X, A) is a cofibered pair, then $(X \times 0) \cup (A \times I)$ is a strong deformation retract of $X \times I$.*

PROOF. Let $i: (X \times 0) \cup (A \times I) \subset X \times I$ be the inclusion map, and let

$$r: X \times I \rightarrow (X \times 0) \cup (A \times I)$$

be a retraction. A homotopy

$$D: ir \simeq 1_{X \times I} \text{ rel } (X \times 0) \cup (A \times I)$$

is given by

$$D(x, t, t') = (pr_1 r(x, (1 - t')t), (1 - t')pr_2 r(x, t) + t't) .$$

THEOREM 3. *Suppose that $p: E \rightarrow B$ is a fibration, that A is a strong deformation retract of X , and that there exists a map $\varphi: X \rightarrow I$ such that $A = \varphi^{-1}(0)$. Then any commutative diagram*

$$\begin{array}{ccc} A & \xrightarrow{f''} & E \\ i \cap & & \downarrow p \\ X & \xrightarrow{f'} & B \end{array}$$

may be filled in with a map $f: X \rightarrow E$ such that $pf = f'$ and $fi = f''$. f is unique up to homotopy rel A .

PROOF. By hypothesis there exists a retraction $r: X \rightarrow A$ and a homotopy

$$D: ir \simeq 1_X \text{ rel } A .$$

If $f: X \rightarrow E$ is such that $fi = f''$, then $f \simeq fir = f''r \text{ rel } A$, which proves the last assertion of the theorem. Define $\bar{D}: X \times I \rightarrow X$ by

$$\bar{D}(x, t) = \begin{cases} D(x, t/(\varphi(x))), & t < \varphi(x), \\ D(x, 1), & t \geq \varphi(x). \end{cases}$$

D is easily shown to be continuous. Because p is a fibration there exists a homotopy $\bar{F}: X \times I \rightarrow E$ such that $p\bar{F} = f'\bar{D}$ and $\bar{F}(x, 0) = f''r(x)$ for each $x \in X$. f is given by $f(x) = \bar{F}(x, \varphi(x))$.

We are now in a position to prove

THEOREM 4. *Suppose that $p: E \rightarrow B$ is a fibration, that (X, A) is a cofibered pair, and that A is closed. Then any commutative diagram*

$$\begin{array}{ccc} (X \times 0) \cup (A \times I) & \xrightarrow{f} & E \\ \cap & & \downarrow p \\ X \times I & \xrightarrow{F} & B \end{array}$$

may be filled in with a homotopy $\bar{F}: X \times I \rightarrow E$ such that $p\bar{F} = F$ and $\bar{F}|_{(X \times 0) \cup (A \times I)} = f$.

PROOF. According to the Lemma $(X \times 0) \cup (A \times I)$ is a strong deformation retract of $X \times I$, and by Theorem 2 there exists a function $\psi: X \rightarrow I$ such that $A = \psi^{-1}(0)$. Define $\varphi: X \times I \rightarrow I$ by $\varphi(x, t) = t\psi(x)$. Then $(X \times 0) \cup (A \times I) = \varphi^{-1}(0)$, and the theorem follows from Theorem 3.

The condition that A be closed is not very restrictive. For instance, A will always be closed if X is Hausdorff. Not all cofibrations are closed, however. The most trivial example of a non-closed cofibration is the pair (X, a) where X is the two-point space $\{a, b\}$ with the trivial topology.

REFERENCES

1. C. H. Dowker, *Homotopy extension theorems*, Proc. London Math. Soc. (3) 6 (1956), 100–116.
2. G. S. Young, *A condition for the absolute homotopy extension property*, Amer. Math. Monthly 71 (1964), 896–897.